

STRONG MGII ABSORPTION SYSTEMS IN QSOS FROM THE SLOAN DIGITAL SKY SURVEY EARLY DATA RELEASE.

Daniel B. Nestor, Sandhya Rao, David Turnshek, Eric Furst*

University of Pittsburgh

dbn@phyast.pitt.edu

1. Introduction

The Sloan Digital Sky Survey (SDSS) is obtaining multicolor images over more than 10,000 square degrees of high Galactic latitude sky and providing medium resolution spectra for approximately 10^6 galaxies and 100,000 quasars. The early data release (EDR) of June 2001 (Schneider et al. 2002) contains spectra of ≈ 3800 QSOs with redshifts ranging from $z = 0.15$ to $z = 5.03$. The spectra cover the wavelength interval $3800\text{\AA} < \lambda < 9200\text{\AA}$ and have resolutions ranging from 1800 to 2100.

In order to study intervening low-ionization metal absorption-line systems, we are constructing samples of systems selected for the strength of their $\lambda 2796$ MgII lines from the EDR. There are 640 absorption systems in our unbiased sample with $\lambda 2796$ REW > 1.0 $\text{\AA}$ detected at 5σ and with redshifts $0.37 < z < 2.27$. This sample is an order of magnitude larger than previous similarly selected samples. In this contribution, we present preliminary results on the statistical properties of the sample.

References

Schneider, et al. 2002, AJ, 123, 567

*also Bucknell University




Figure 1 **Left:** The redshift path covered by the sample. **Right:** The number density distribution with redshift of the sample. Also shown are no-evolution predictions for three cosmologies normalized to minimize the χ^2 fit to the binned data. The distribution is consistent with no evolution except at the lowest redshifts.




Figure 2 **Left:** Evolution of the REW distribution slope with redshift for $n(W_0) = n_0 e^{-W_0/W_0^*}$. The data are shown divided into redshift bins indicated by horizontal bars with an equal number of lines in each bin. Vertical bars are 1σ errors. There is mild evolution in the distribution of REWs. **Right:** The REW distribution for the total sample. The slope of the maximum-likelihood fit is -1.4 , corresponding to $W_0^* = 0.7$.